

# 2-[2-(4-Ethoxyphenyl)methyl-5-nitro-1*H*-benzimidazolyl]-*N,N*-diethylethanaminium Chloride-Acetic Acid: Etonitazene

By C. HUMBLET, G. EVRARD AND F. DURANT

Laboratoire de Chimie Moléculaire Structurale, Groupe de Chimie-Physique, Facultés Universitaires Notre-Dame de la Paix, 61 rue de Bruxelles, B-5000 Namur, Belgium

(Received 29 July 1978; accepted 1 September 1978)

**Abstract.**  $C_{22}H_{28}N_4O_3 \cdot HCl \cdot C_2H_4O_2$ ,  $P2_1/c$ ,  $a = 9.721(5)$ ,  $b = 13.588(5)$ ,  $c = 19.673(5) \text{ \AA}$ ,  $\beta = 92.47(5)^\circ$ ,  $D_m = 1.25$ ,  $D_c = 1.26 \text{ g cm}^{-3}$ ,  $Z = 4$ . The planes of the benzimidazole groups are stacked parallel.

**Introduction.** The present study was undertaken as part of an investigation of the structure-activity relationship in narcotic analgesics. The title compound is a thousand times more potent than morphine.

Colourless prismatic crystals were obtained from ethyl acetate. The hydrochloride form of the starting compound could explain the solvent hydrolysis which gave rise to the acetic acid found in the crystal structure.

The space group was determined from photographs. Cell dimensions and intensities were measured on a Nonius CAD-4 diffractometer with the experimental conditions given in Table 1. The structure was solved with MULTAN (Germain, Main & Woolfson, 1971). The most probable *E* map showed all atoms except H.

Full-matrix least-squares refinement with XRAY 72 (Stewart, Kruger, Ammon, Dickinson & Hall, 1972) converged at  $R = 0.14$ . A difference map revealed disorder at the end C of one of the *N*-ethyl chains; the two positions selected and the corresponding population factors were included in the refinement. The H atoms of the two hydrogen bonds were observed on this map. Refinement with anisotropic temperature factors for Cl, O, N and C and isotropic for H converged at a final  $R = 0.061$  (for observed

Table 1. Experimental conditions

Source: Cu  $K\bar{\alpha}$ ,  $\lambda = 1.54178 \text{ \AA}$   
 Scan:  $\omega=2^\circ$   
 Graphite monochromator  
 Confidence level:  $2.5\sigma$ , with  $\sigma^2(I) = S + B + (0.03S)^2$  ( $S$  being the scan and  $B$  the background count)  
 $2.0 \leq \theta \leq 70.0^\circ$   
 Scanning angle:  $0.8 + 0.2 \tan \theta (\text{deg})$   
 Aperture :  $2.5 + 0.5 \tan \theta (\text{mm})$   
 Total number of independent reflexions: 3282  
 Number observed: 2236

reflexions). The final positional parameters are given in Table 2.\*


\* Lists of structure factors and anisotropic thermal parameters have been deposited with the British Library Lending Division as Supplementary Publication No. SUP 33852 (32 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

Table 2. Final positional parameters ( $\times 10^4$ ) with e.s.d.'s in parentheses

	x	y	z
C(1)	-186 (11)	1383 (8)	920 (7)
C(2)	1075 (11)	1866 (8)	1269 (6)
O(3)	1863 (5)	1046 (4)	1546 (3)
C(4)	3063 (7)	1247 (5)	1911 (4)
C(5)	3755 (7)	435 (5)	2196 (4)
C(6)	4967 (7)	568 (5)	2580 (3)
C(7)	5503 (6)	1505 (5)	2698 (3)
C(8)	4811 (7)	2300 (5)	2406 (4)
C(9)	3598 (7)	2191 (5)	2013 (4)
C(10)	6804 (6)	1650 (5)	3150 (3)
C(11)	6618 (6)	2334 (5)	3734 (3)
N(12)	7271 (5)	3178 (4)	3806 (3)
C(13)	6839 (6)	3567 (5)	4414 (3)
C(14)	7196 (6)	4454 (5)	4731 (3)
C(15)	6605 (7)	4619 (5)	5339 (3)
N(16)	6911 (7)	5565 (5)	5685 (4)
O(17)	7717 (7)	6121 (4)	5418 (3)
O(18)	6319 (7)	5763 (4)	6204 (3)
C(19)	5693 (7)	3990 (5)	5645 (3)
C(20)	5319 (7)	3108 (5)	5326 (3)
C(21)	5907 (6)	2927 (5)	4700 (3)
N(22)	5767 (5)	2147 (4)	4252 (2)
C(23)	4875 (6)	1293 (5)	4335 (3)
C(24)	3420 (6)	1525 (5)	4061 (3)
N(25)	2420 (5)	752 (4)	4261 (3)
C(26)	938 (8)	1048 (7)	4092 (5)
C(27)	409 (14)	1838 (12)	4420 (9)
C(27)	671 (18)	1392 (17)	3421 (12)
C(28)	2746 (8)	-242 (6)	3969 (4)
C(29)	1923 (10)	-1059 (6)	4280 (5)
Cl(60)	2785 (2)	691 (1)	5798 (1)
C(50)	11170 (9)	4369 (8)	2528 (4)
C(51)	9841 (9)	3914 (7)	2720 (4)
O(52)	9671 (5)	3877 (4)	3352 (3)
O(53)	8950 (7)	3660 (7)	2318 (3)

**Discussion.** The atomic numbering, bond distances and angles are given in Fig. 1. The main torsion angles are listed in Table 3. Fig. 2 is a stereoview of the molecule.

There are two strong hydrogen bonds from the analgesic molecule to the  $Cl^-$  ion and acetic acid molecule:  $Cl(60) \cdots N(25) = 3.034$ ,  $N(25)-H(601) = 1.002$ ,  $Cl(60) \cdots H(601) = 2.036 \text{ \AA}$ ,  $Cl(60) \cdots H(601)-N(25) = 173.49^\circ$ ,  $O(52) \cdots N(12) = 2.692$ ,  $O(52)-H(521) = 1.152$ ,  $N(12) \cdots H(521) = 1.546 \text{ \AA}$ ,  $O(52)-H(521) \cdots N(12) = 172.23^\circ$ .



(a)




Fig. 1. (a) Bond distances ( $\text{\AA}$ ) and (b) angles ( $^\circ$ ). The average e.s.d.'s for the lengths are  $0.009 \text{ \AA}$  and for angles  $0.6^\circ$  except for the terminal ethyl groups for which e.s.d.'s range between  $0.026-0.012 \text{ \AA}$  and  $0.8-1.3^\circ$ .




Fig. 2. Stereoscopic view (Johnson, 1965) of the molecule with 50% probability thermal ellipsoids.

Table 3. Main torsion angles ( $^\circ$ )

C(6)-C(7)-C(10)-C(11)	-123.5
C(8)-C(7)-C(10)-C(11)	55.2
C(7)-C(10)-C(11)-N(22)	-116.2
C(10)-C(11)-N(22)-C(23)	-1.5
C(11)-N(22)-C(23)-C(24)	-95.5
N(22)-C(23)-C(24)-N(25)	-158.6
C(23)-C(24)-N(25)-C(26)	170.8
C(23)-C(24)-N(25)-C(28)	-62.6
C(24)-N(25)-C(28)-C(29)	168.5
C(24)-N(25)-C(26)-C(27)	-64.8
C(1)-C(2)-O(3)-C(4)	-178.4
C(2)-O(3)-C(4)-C(5)	177.0
C(2)-O(3)-C(4)-C(9)	-2.3
C(14)-C(15)-N(16)-O(17)	-2.9
C(19)-C(15)-N(16)-O(18)	-2.5




Fig. 3. Stereoscopic view of the packing.

The packing is essentially due to van der Waals interactions between the almost parallel benzimidazole groups of the  $(x,y,z)$  and  $(1-x, 1-y, 1-z)$  equivalent molecules, the interplanar distances ranging from  $3.4$  to  $3.5 \text{ \AA}$ . The resulting shape of the packing is a pleated sheet (Fig. 3).

We thank the Ciba-Geigy pharmaceutical laboratories and Dr J. M. Tyberghein (Ciba-Geigy, B-1720 Groot-Bijgaarden, Belgium) for providing the sample and the Institut pour l'Encouragement de la Recherche Scientifique dans l'Industrie et l'Agriculture (IRSIA) for the award of a doctoral fellowship to CH.

## References

- GERMAIN, G., MAIN, P. & WOOLFSON, M. M. (1971). *Acta Cryst. A* 27, 368–376.
- JOHNSON, C. K. (1965). ORTEP. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee.
- STEWART, J. M., KRUGER, G. J., AMMON, H. L., DICKINSON, C. & HALL, S. R. (1972). The XRAY system – version of June 1972. Tech. Rep. TR-192. Computer Science Center, Univ. of Maryland, College Park, Maryland.