

# Human Variation in Pelvic Shape and the Effects of Climate and Past Population History

LIA BETTI\*

Centre for Research in Evolutionary, Social and Inter-Disciplinary Anthropology,  
Department of Life Sciences, University of Roehampton, London, SW15 4JD, UK

---

---

## ABSTRACT

The human pelvis is often described as an evolutionary compromise (obstetrical dilemma) between the requirements of efficient bipedal locomotion and safe parturition of a highly encephalized neonate, that has led to a tight fit between the birth canal and the head and body of the foetus. Strong evolutionary constraints on the shape of the pelvis can be expected under this scenario. On the other hand, several studies have found a significant level of pelvic variation within and between human populations, a fact that seems to contradict such expectations. The advantages of a narrow pelvis for locomotion have recently been challenged, suggesting that the tight cephalo-pelvic fit might not stem from the hypothesized obstetrical dilemma. Moreover, the human pelvis appears to be under lower constraints and to have relatively higher evolvability than other closely related primates. These recent findings substantially change the way in which we interpret variation in the human pelvis, and help make sense of the high diversity in pelvic shape observed within and among modern populations. A lower magnitude of covariance between functionally important regions ensured that a wide range of morphological variation was available within populations, enabling natural selection to generate pelvic variation between populations living in different environments. Neutral processes such as genetic drift and differential migration also contributed to shaping modern pelvic diversity during and after the expansion of humans into and across the various continents. *Anat Rec*, 300:687–697, 2017. © 2017 Wiley Periodicals, Inc.

**Key words:** pelvis; climate; evolvability

---

---

The last few years have seen a remarkable increase in research on the evolution of the human pelvis and on the factors that explain pelvic shape variation in modern human populations. The classic model of the human pelvis as a strict compromise between locomotory and obstetrical requirements, omnipresent in anthropology textbooks, has been challenged from different directions, first of all by calling into question the underlying idea of a conflict between a wide, obstetrically capacious pelvis and efficient bipedal locomotion. The accompanying assumption of strong evolutionary constraints on the shape of the pelvis has also been disputed, and new evidence has emerged of ample geographic variation in pelvic shape within our species. While geographic variation in the pelvis has often been explained as climatic

adaptation, new research has revealed the overarching effects of neutral evolutionary processes in generating population differences in pelvic shape. This review integrates old and new evidence to draw a sketch of the

---

\*Correspondence to: Lia Betti, Centre for Research in Evolutionary, Social and Inter-Disciplinary Anthropology, Department of Life Sciences, University of Roehampton, London, SW15 4JD, UK. E-mail: Lia.Betti@roehampton.ac.uk

Received 15 March 2016; Revised 30 August 2016; Accepted 20 October 2016.

DOI 10.1002/ar.23542

Published online in Wiley Online Library (wileyonlinelibrary.com).




Fig. 1. Caldwell and Moloy's pelvic types (Caldwell and Moloy, 1938; from an unknown original source).


current state of understanding of the origins of human pelvic shape variation.

### Early Observations of Pelvic Variation and the "Obstetrical Dilemma" Hypothesis

Humans originated in Africa, but have dispersed globally and colonised a variety of environments. As a consequence, modern human populations show a genetic and phenotypic variation that is geographically structured. For centuries, and up to the middle of the 20th century, many naturalists and anthropologists made sense of human variation by assigning populations to well-defined groups based on their physical and cultural characteristics. Early studies of human pelvic variation followed this approach, and focused on the prevalence of different pelvic shapes (e.g., Fig. 1) in the various 'races'

as well as their obstetric proficiency (e.g., Turner, 1885; Caldwell and Moloy, 1933; Greulich and Thoms, 1938; Allen, 1947a,b). The rounder shape of the canal in some populations (e.g., KhoeSan, Andamanese, and Australians) was considered simultaneously to be more primitive and more suitable for childbirth. The transversally more elongated shape of European women, on the other hand, was described as less adept for childbirth being the race farthest from the animal kingdom (Turner, 1885—who was, incidentally, European). Although the female pelvis was generally assumed to be shaped in accordance to its obstetric functions, it became obvious that it varied in shape between human groups.

Since the mid 20th century, most scholars moved away from a racial categorization of human variation, and pelvic shape became largely discussed as the outcome of evolutionary processes. The human pelvis differs



The basic tenet of the obstetrical dilemma is that a narrower bi-acetabular distance, which positions the support of the body closer to the centre of gravity, reduces the rotation of the hip during locomotion and the energy required by the muscles to counteract it, thereby improving efficiency. At the same time, a narrower bi-acetabular breadth would also lead to a less spacious pelvic canal, with an increased risk of foeto-pelvic disproportion and obstructed labour, where the head or body of the neonate does not fit through the mother's birth canal (Rosenberg, 1992; Wittman and Wall, 2007). The selective pressure for efficient locomotion and the female-specific pressure for a sufficiently large pelvic canal led to a compromise, whereby the female pelvis is larger than the male pelvis, to accommodate the passage of the foetus, but probably less efficient for locomotion. The result of this compromise is a very tight fit between the newborn's head and shoulders and the mother's birth canal, balancing the two contrasting needs (Schultz, 1949; Leutenegger, 1974; Tague and Lovejoy, 1986; Rosenberg, 1992; Rosenberg and Trevathan, 1995; Trevathan and Rosenberg, 2000; DeSilva, 2011). Adaptations to bipedalism are also considered a likely explanation for the twisted human birth canal,

A study by Huseynov et al. (2016) recently suggested that evolutionary constraints on the increase of the distance between the ischial spines, which would alleviate foeto-pelvic disproportion, might be due to the need to provide support for abdominal organs instead of increased locomotory efficiency. In fact, important ligaments and fascias that form the pelvic floor are anchored to the ischial spines (Abitbol, 1988). The authors also highlighted how pelvic shape changes

during an individual's life, whereby canal dimensions reach a peak during the female most fertile years (around 20–40 years of age), followed by a slight decrease in canal size at the level of the ischial spines in the postmenopausal period. These female-specific shape changes over the life cycle might indicate long-term adaptation to ensure obstetric sufficiency during fertile years.

Obstetric sufficiency can, indeed, be expected to be under strong selective pressure. Complications during childbirth can lead to the death of the newborn and of the mother, or cause damage that could make future births difficult. As such, the impact on the mother's fitness is potentially extremely high. Considering that childbirth mortality and morbidity are not insignificant in modern populations, especially when hospital care is not available (Kolawole et al., 1978; Oxorn, 1986; Cottingham and Royston, 1991; Abou Zahr and Wardlaw, 2004; Dolea and AbouZhar, 2003; Hogan et al., 2010), it is surprising that our species has not evolved a larger birth canal.

Grabowski (2013) hypothesized that a possible reason for the tight fit observable in modern women could be that there has simply not been enough time for the pelvis to adapt to the rapid increase in encephalization that occurred in Middle Pleistocene hominins (Ruff et al., 1997; Rightmire, 2004), possibly due to a higher level of genetic constraints on the pelvis in respect to neonatal cranial size (Grabowski, 2013).

Another possible explanation is that the tight fit observed today is a relatively recent consequence of a change in diet and environmental conditions. Differential phenotypic plasticity (i.e., the capacity for phenotypic changes in response to environmental changes) in the pelvis and in neonatal size might mean that the likelihood of foeto-pelvic disproportion varies depending on contingent environmental condition. Wells et al. (2012) suggest that the fit between neonate and canal became tighter after the development of agriculture. The decreased protein intake and increased glycemic load associated with the change in subsistence would have resulted in lower adult stature of the mother but higher neonatal weight, therefore exasperating what was originally a close, but not as tight, fit. The recent shift towards a highly calorific diet could have exasperated further the situation. If plasticity is at the root of the problem, the obstetrical dilemma might not be a dilemma after all. The size of the canal evolved to be sufficient in previous environmental conditions, and selection is now catching up with the change in circumstances.

Kurki (2007) noted that the KhoeSan population of South Africa, who tend to have a relatively small body size, appear to have a larger-than-expected birth canal when compared to bigger-bodied populations. The dimensions involved have obstetric significance, indicating that the selective pressure for a spacious birth canal could be particularly strong in small-bodied populations, possibly because the head of the newborn is not isometrically scaled according to the mother's body size, and is relatively larger than in bigger-bodied populations (Wells et al., 2012). The size of the canal might be protected in human populations, despite its large variability in shape (Kurki, 2013b). Indeed, the foetal head is quite plastic, due to the fact that the cranial bones are not fused and to the presence of the fontanelles, and it can

partially adapt its shape while negotiating the mother's birth canal (e.g., Stewart and Philpott, 1980; Kriewall and McPherson, 1981).

Shorter-than-average mothers appear to have a higher probability of birth complications in various human populations (e.g., Tsu, 1992; Kwawukume et al., 1993; Witter et al., 1995; Cnattingius et al., 1998; Prasad and Al-Taher, 2002; Sheiner et al., 2005) due to a more frequent mismatch between the size of the neonate and the size of the pelvic canal. The probability of a mismatch is reduced in part by the fact that neonatal head size tends to be inherited from the mother, so that small mothers with a small head tend to give birth to small-headed babies (Leary et al., 2006). In addition, females with a larger head, who are more likely to give birth to babies with a larger head, tend to have a rounder, obstetrically more capable birth canal (Fischer and Mitteroecker, 2015). A rounder birth canal is also more likely in shorter women, who are at higher risk of cephalo-pelvic disproportion (Fischer and Mitteroecker, 2015). This set of correlations indicates long-term selective pressure due to obstetric requirements.

Despite some evidence of obstetric-related selection, the ample variation in the shape of the pelvis and of the birth canal in modern women, both within and among populations, brings into question the obstetrical dilemma as a strict compromise between locomotion efficiency and obstetric sufficiency. If strong evolutionary constraints due to a shared selective pressure (i.e., bipedal locomotion) were acting on the pelvis, we would expect to find low variability in the shape of the pelvic structure; however, not only does the birth canal shows a higher shape variation than other areas of the pelvis (Kurki, 2013b), but women are not less variable than men in this respect (Tague, 1989, 1995; Arsuaga and Carretero, 1994; LaVelle, 1995; Betti et al., 2013; Kurki, 2013b).

### Human Populations Show Ecogeographic Variation in Pelvic Shape

Beyond early studies of racial variation in the pelvis, which highlighted differences between races in the shape of the pelvic canal (e.g., Turner, 1886; Caldwell and Moloy, 1933) and of the ilium (Derry, 1923; Straus, 1927), more recent studies have confirmed significant variation in pelvic shape between populations. Işcan (1983) and Patriquin et al. (2002) found that it was possible to set apart individuals of 'white' and 'black' ancestry based on a set of pelvic measurements with an accuracy of over 80%. Weaver and colleagues examined variation in the whole articulated pelvis in several human populations using 3D landmark coordinates, and were able to show that modern populations vary significantly in overall pelvic shape (Weaver, 2002; Weaver and Hublin, 2009). Similar results were obtained by Betti et al. (2014), who focused on the 3D shape of the ilium in a large number of populations. Wells et al. (2012) compared several studies on skeletal and in vivo pelvic canal measurements and were able to show that high variation between populations is also present in the birth canal.

Variation in pelvic breadth among populations, in particular, has been noted and discussed by several authors (e.g., Ruff, 1991, 1993, 1994; Holliday and Hilton, 2010,



Kurki, 2013a). Populations from higher latitudes tend to have relatively wider bi-iliac breadth than populations from lower latitudes, suggesting that part of the variation might be related to climate. There is abundant evidence that populations living in different climates tend to have different body proportions, with cold-adapted populations having relatively larger and stockier bodies, and populations in hot climates having smaller or more elongated bodies with relatively longer limbs (Schreider, 1950, 1964, 1975; Newman, 1953, 1961; Roberts, 1953, 1978; Baker, 1960; Coon, 1962; Crognier, 1981; Ruff, 1994, 2002; Holliday, 1997a,b, 1999, 2002; Katzmarzyk and Leonard, 1998; Weaver, 2002; Weaver and Steudel-Numbers, 2005; Weinstein, 2005; Tilkens et al., 2007; Weaver and Hublin, 2009; Betti et al., 2012, 2015). This pattern is widespread in endotherm animals, with closely related species displaying larger body sizes and shorter extremities in colder environments, and it is often referred to as Bergmann's (1847) Rule (for body size) and Allen's (1877) Rule (for body and limb proportions).

*Homo sapiens* evolved in Africa, where most regions are characterised by a warm or hot climate. For endothermic species such as humans, maintaining a stable body temperature in hot environments requires efficient heat dissipation; this is achieved in our species through sweat evaporation at the body surface, which is facilitated by a high body surface-to-volume ratio. The relatively elongated body shape of most sub-Saharan African populations in respect to other human populations, and the associated higher surface-to-volume ratio, has been interpreted as a thermoregulatory adaptation to a warm environment (Ruff, 1994, 2002; Holliday and Falsetti, 1995; Holliday, 1997a; Holliday and Hilton, 2010). After the expansion out of Africa and the colonisation of high latitude regions, it has been suggested that natural selection favoured more robust and stocky bodies with a lower surface-to-volume ratio, which helped maintain internal body temperature in cold climates. Differential climatic adaptation might therefore have contributed in generating modern population differences in body proportions, body size, limb robusticity, and even pelvic shape (Trinkaus, 1981; Ruff, 1991, 1994; Holliday and Falsetti, 1995; Pearson, 2000; Stock, 2006; Weaver and Hublin, 2009).

Kurki (2013a) compared nine populations from various continents, and found a relationship between latitude and canal shape, whereby high-latitude populations tend to have a transversally wider midplane and outlet and an inlet which is less deep antero-posteriorly than mid- and low-latitude populations. Betti et al. (2014) and Weaver (2002) found a significant relationship between temperature and the 3D shape of the ilium and of the whole pelvis, suggesting that both pelvic and canal shape have been partially shaped by local climatic adaptations, possibly through the effect of natural selection favouring thermally efficient body sizes and proportions.

It is possible, and indeed it has been suggested before (e.g., Ruff, 1994; Weaver and Hublin, 2009; Kurki, 2013a), that thermoregulatory adaptations in the shape of the pelvis generated a different type of conflict with obstetric requirements: another form of 'dilemma'. The narrower body favoured in hot and humid environments could limit the transversal width of the birth canal, increasing at the same time the anterior-posterior diameter in order to preserve a minimum size for the passage

of the foetus; on the contrary, the wider body common in cold regions could translate into a more transversally elongated shape of the canal at higher latitudes. This interpretation could help make sense of the geographic variation in the pelvic canal recognised since early racial studies, taking into account, at the same time, shared obstetric requirements and constraints.

In addition to climate, several other factors are thought to contribute in generating pelvic shape variation within and between human populations. For example, nutritional deficiencies and disease, sometimes associated with low social status, as well as activity, age and number of pregnancies, have been shown to have an influence on skeletal and pelvic size and shape (Greulich et al., 1939; Nicholson, 1946; Thoms, 1946, 1947; Krukierek, 1951, 1954; Bernard, 1952; Angel, 1978; Kelley, 1978; Holland et al., 1982; Micozzi, 1982; Abitbol, 1987; Kelley and Angel, 1987; Stuart-Macadam, 1989; Sibley et al., 1992; Bogin et al., 2002; Bogin and Rios, 2003; Specker and Binkley, 2005; Schutz et al., 2009; Huseynov et al., 2016). However, these factors are expected to explain differences between individuals within a population more than differences between populations, except in case of substantial and systematic differences in diet, lifestyle or health between certain populations.

### **Evolvability of the Pelvis and the Effects of Past Population History**

In a comparative study of the primate pelvic girdle, Grabowski (2013) examined the pattern and strength of covariation between pelvic measurements in different species. He found a significantly smaller covariance in the dimensions of the human pelvis compared with other closely related species. This result is surprising in the light of the fact that humans experience a tighter cephalo-pelvic fit during childbirth in respect to other apes (Abitbol, 1991; Rosenberg and Trevathan, 2002), a fact that should hypothetically translate into stronger evolutionary constraints on the size and shape of the female pelvic canal. Grabowski and colleagues (Grabowski et al., 2011; Grabowski, 2013) interpret the lower human covariance as the outcome of the shift to bipedalism, whereby a release of constraints allowed for a substantial rearrangement of the shape of the hominin pelvis in australopithecine or earlier species. The decrease in covariance comes with an increase in the evolvability of the pelvic structure, which is freed from previous constraints and more susceptible to change (Grabowski, 2013). Lower covariance also means that it is relatively easier for a structure to evolve independently of other structures, so that different sets of traits could evolve in response to different selective pressures.

Middle Pleistocene hominins showed a rapid increase in encephalization from around 500 ky years ago (Ruff et al., 1997; Rightmire, 2004), leading to a tight cephalo-pelvic fit in humans and Neanderthals. Obstetric requirements were fulfilled in different ways in the two later Pleistocene species, probably taking advantage of the relatively high pelvic evolvability. Neanderthal females achieved a large pelvic canal thanks to a more elongated pubic ramus (Rosenberg et al., 1988). Neanderthal and modern human birth canals differ also in the geometry of the three pelvic planes. Although different reconstructions of the only fairly complete




Fig. 3. Plots of between-population phenotypic distance against geographic distance for the cranium (A) and the os coxae (B). Data from Betti et al., 2010, 2014.

Neanderthal female pelvis, Tabun C1, have been put forward, it seems likely that Neanderthals preserved the ancestral transverse orientation of the main diameter of the outlet, therefore presenting a different pattern of childbirth from modern humans (Weaver and Hublin, 2009; but see Ponce de León et al., 2008). The changing pattern of covariance between pelvic traits and the likelihood of indirect effects of selection on covariant traits over hominin evolution underlines the importance of using a multivariate approach to understanding the evolution of pelvic morphology (Rolian, 2014; Grabowski and Roseman, 2015; Grabowski, 2016).

A lower level of covariance and a relatively higher evolvability in the human pelvis could help explain the large shape variation observed within and among modern populations. The lack of strong evolutionary constraints on pelvic shape helps make sense of why, against expectations due to obstetric selective pressures, the pelvic canal shows high variability, and that the level of variation is the same in males and females.

Low constraints on the pelvis can also explain the clear ecogeographic pattern in pelvic shape observed in human populations, despite the fact that their recent dispersal into colder environments offered only a short time for evolution to act. The greater freedom of pelvic traits to change independently on each other offers a wider range of possible shapes that can evolve quickly without disrupting the whole pelvic structure. This freedom returned a high diversity within-population, which is the ideal basis for directional selection to improve adaptation to different environments.

A higher magnitude of pelvic evolvability also offers a wider scope for genetic drift to increase phenotypic

differences between populations. Our species is relatively young, probably not much older than 200–250 ky old, and went through a rapid range expansion of a global scale in the last 100–60 ky (e.g., McDougall et al., 2005; Mellars, 2006; Armitage et al., 2011; Eriksson et al., 2012; Fu et al., 2013). The rapid geographic expansion was accompanied by a reduction in genetic diversity with increasing distance from the continent of origin, Africa (Prugnolle et al., 2005; Ramachandran et al., 2005). This loss of diversity is evident in both genetic and phenotypic variation, such as cranial and dental shape variation (Manica et al., 2007; Hanihara, 2008; von Cramon-Taubadel and Lycett, 2008; Betti et al., 2009), and is the natural consequence of genetic drift when a series of founder events occur. Each subpopulation which colonised a new area carried only a portion of the genetic diversity present in the source population, thereby generating the described pattern of gradual decrease of genetic variation.

A similar decline in phenotypic variation with distance from Africa has been identified in the os coxae (Betti et al., 2013) (Fig. 3). The signature of the species' expansion appears to be even stronger on the pelvis than on the cranium, with distance from sub-Saharan Africa explaining up to about 47% of within-population variance (Betti et al., 2013), a fact that indicates that modern variation in the os coxae has been significantly affected by genetic drift.

As human range expansion followed divergent routes into the various continents, it might be expected that genetic drift led to a random shift in average pelvic shape along the main directions of expansion. In fact, shape differences in the os coxae among populations appear to increase with geographic distance along those different routes (Betti et al., 2014) (Fig. 4). The colonisation of new areas and subsequent exchange of migrants between populations are expected to generate a geographic pattern of variation, with nearby populations showing closer genetic and phenotypic resemblance than populations more distant in space. This pattern is due to the fact that closer populations are more likely to have emerged from a recent common ancestral population during the expansion of our species; in addition, geographic proximity favours gene flow between populations and increases similarity (Imaizumi et al., 1973; Cavalli-Sforza et al., 1994; Eller, 1999; Relethford, 2004b). Both the cranium and the pelvis show a similar pattern, with geographic distance explaining about 30–35% of between-population shape variance (Betti et al., 2010, 2014; see also Roseman, 2004; Harvati and Weaver, 2006a; Hubbe et al., 2009; von Cramon-Taubadel, 2009a) (Fig. 4).

In comparison, climatic adaptation plays a significant, but smaller role. Differences in climate explain up to about 18% of coxal shape variation among populations, which is reduced even further once the variation explainable by neutral processes is taken into account (Betti et al., 2014). Climate, on the other hand, seems to have a stronger effect on coxal size, with minimum temperature explaining up to 43% of size variation (Betti et al., 2014).

Pelvic sexual dimorphism also appears to have been affected by past demographic events (Betti, 2014). Shape differences between the sexes are not stable across human populations, but vary geographically. Part of this geographic diversity in sexual dimorphism (ca. 15%;




Fig. 4. Plots of within-population phenotypic variance against geographic distance from sub-Saharan Africa for the cranium (A) and the os coxae (B). Data from Betti et al., 2009, 2013.

Betti, 2014) can be explained, again, by population history.

Although the amounts of variance explained by neutral factors need to be considered as purely indicative, stochastic processes due to genetic drift and differential migration clearly played an important role in shaping pelvic differences between populations. Climatic adaptation acted over this underlying neutral pattern, affecting both pelvic size and shape. The comparison between the effects of climatic adaptation and past population history, however, suggest that the latter had a larger influence on pelvic—or at least, coxal—shape variation. The importance of neutral processes is also reflected in the global pattern of apportionment of pelvic shape variance. As for neutral genetic markers and cranial shape, pelvic shape diversity between populations and between continents is much smaller than the average diversity within single populations (Betti et al., 2013). The opposite is true for skin colour, for example, whereby strong directional selection increased the variance among regions well beyond the variance within populations (Relethford, 2002).

Acknowledging that a substantial portion of phenotypic variance in modern populations is the outcome of neutral processes is important for at least two reasons. First, it provides an explanation for morphological differences which are not clearly related to local adaptations. Second, it affects the way we test for natural selection. Adaptation due to natural selection is often inferred from a significant correlation between the factor of interest and morphology. However, environmental factors are usually partially correlated with geographic distance, due to the fact that nearby areas tend to share similar environmental conditions. As morphological distance also tends to be correlated with geographic distance,

because of neutral demographic processes, a significant correlation between morphological traits and environmental factors could be due, at least in part, to collinearity with geographic distance. In other words, the underlying neutral pattern of variation could confound the analyses and generate spurious correlations with environmental factors. Roseman and Auerbach (2015) showed very eloquently how this problem can impact on tests of climate-related selection on the human postcranium.

## CONCLUSIONS

The view of the pelvic girdle as a strict compromise between two opposite selective pressures, a narrow bi-acetabular distance for efficient bipedal locomotion, and a wide pelvic canal for easy parturition, has been challenged (Dunsworth et al., 2012; Wall-Scheffler, 2012; Warrener et al., 2015). The wider female pelvic breadth is not associated with less efficient locomotion, calling into question the elegant hypothesis of an “obstetrical dilemma.” This finding leaves open the question of why a wider female pelvic canal has not evolved in our species, given the significant risk of childbirth mortality and morbidity due to photo-pelvic disproportion and the subsequent decrease in fitness. One possible explanation is that the modern sedentary lifestyle and the availability of highly calorific food enjoyed by many populations have exacerbated the risk of childbirth complications by increasing average neonatal size (Wells et al., 2012). If that were the case, the hypothesized selective advantage of a larger birth canal would be a recent phenomenon.

These studies not only cast doubt on the existence of an obstetrical dilemma as it was originally framed, but also on the existence of strong evolutionary constraints on the pelvic structure, which would be the logical extension of the obstetrical dilemma hypothesis. Indeed, the pelvis and the pelvic canal do not show low morphological variation in respect to other parts of the skeleton, as would be expected under strong evolutionary constraints (Kurki, 2007, 2013a,b; Betti et al., 2014). On the contrary, it appears that the human pelvis is under lower constraints than other closely related primates, showing lower covariance between its parts and a higher level of evolvability (Grabowski, 2013; Grabowski et al., 2011).

The evidence of low evolutionary constraints on the pelvis helps make sense of the wide diversity in pelvic and pelvic canal shape observed in modern human populations. A relatively high evolvability means that a larger range of possible shapes can evolve in a shorter time, since it can provide more variation upon which both natural selection and drift can act. The rapid expansion of our species out of Africa into higher latitudes and colder environments exposed human populations to new selective pressures, which could have contributed to the geographic variation in pelvic shape observable today. In fact, the wider pelvis of high-latitude populations and the narrower bodies of tropical populations have been interpreted as a thermoregulatory adaptation (Ruff, 1991, 1993, 1994; Holliday, 1997b; Holliday and Hilton, 2010; Kurki, 2013a). Differences in body size in human populations could also have been accompanied by increased selective pressure for obstetric capacity,



leading to a relatively larger and deeper birth canal in small-bodied populations (Kurki, 2007).

A substantial portion of pelvic variation among modern populations has been shaped by neutral evolutionary processes, such as genetic drift and distance-mediated gene flow (Betti et al., 2014). In the study of human evolution and diversity, there has been a tendency to interpret skeletal variation as the direct effect of natural selection, employing an adaptationist approach, while the effects of neutral processes have been largely ignored. Several recent studies, on the other hand, have highlighted the potential impact of random genetic drift in hominin evolution (e.g., Weaver et al., 2007, 2008; Ackermann and Cheverud, 2004; Grabowski and Roseman, 2015) as well as modern human variation (e.g., Relethford, 1994, 2002, 2004a; Roseman, 2004; Roseman and Weaver, 2004; Harvati and Weaver, 2006; Smith, 2009, 2011; Betti et al., 2009, 2010, 2014; von Cramon-Taubadel, 2009a,b, 2011; von Cramon-Taubadel and Weaver, 2009; Strauss and Hubbe, 2010). Despite that fact that adaptation narratives have an undeniable appeal, the less interesting random accumulation of differences due to genetic drift and differential migration is an ever-occurring process and can potentially explain a large portion of phenotypic variation between human populations.

# LITERATURE CITED

- Abitbol MM. 1987. Obstetrics and posture in pelvic anatomy. *J Hum Evol* 16:243–255.
- Abitbol MM. 1988. Evolution of the ischial spine and of the pelvic floor in the Hominoidea. *Am J Phys Anthropol* 75:53–67.
- Abitbol MM. 1991. Ontogeny and evolution of pelvic diameters in anthropoid primates and in *Australopithecus afarensis* (AL 288-1). *Am J Phys Anthropol* 85:135–148.
- Abou Zahr C, Wardlaw T. 2004. Maternal mortality in 2000: estimates developed by WHO, UNICEF and UNFPA. Geneva: World Health Organization.
- Ackermann RR, Cheverud JM. 2004. Detecting genetic drift versus selection in human evolution. *Proc Natl Acad Sci USA* 101:17946–17951.
- Aiello LC, Dean C. 1990. An introduction to human evolutionary anatomy. London: Academic Press.
- Allen EP. 1947a. Standardised radiological pelvimetry. I. Quantitative aspects. *Brit J Radiol* 20:45–54.
- Allen EP. 1947b. Standardised radiological pelvimetry. II. Interpretation of pelvimetry. *Br J Radiol* 20:205–218.
- Allen JA. 1877. The influence of physical conditions in the genesis of species. *Radical Rev* 1:108–140.
- Angel JL. 1978. Pelvic inlet form: a neglected index of nutritional status. *Yearb Phys Anthropol* 48:378.
- Armitage SJ, Jasim SA, Marks AE, Parker AG, Usik VI, Uerpmann H-P. 2011. The southern route “Out of Africa”: evidence for an early expansion of modern humans into Arabia. *Science* 331:453–456.
- Arsuaga JL, Carretero JM. 1994. Multivariate analysis of the sexual dimorphism of the hip bone in a modern human population and in early hominids. *Am J Phys Anthropol* 93:241–257.
- Baker PT. 1960. Climate, culture, and evolution. *Hum Biol* 32:3–16.
- Bass WM. 1971. Human osteology: a laboratory and field manual of the human skeleton. Columbia: Missouri Archaeological Society.
- Bergmann C. 1847. Über die verhältnisse der warmeoekonomie der thiere zu ihrer grosse. *Göttingen Studien* 1:595–708.
- Bernard RM. 1952. The shape and size of the female pelvis. *Trans Edinburgh Obstet Soc* 59:1–16.
- Betti L. 2014. Sexual dimorphism in the size and shape of the os coxae and the effects of micro-evolutionary processes. *Am J Phys Anthropol* 153:167–177.
- Betti L, Balloux F, Hanihara T, Manica A. 2009. Ancient demography, not climate, explains within-population phenotypic diversity in humans. *Proc R Soc B* 276:809–814.
- Betti L, Balloux F, Hanihara T, Manica A. 2010. The relative role of drift and selection in shaping the human skull. *Am J Phys Anthropol* 141:76–82.
- Betti L, Lycett SJ, von Cramon-Taubadel N, Pearson OM. 2015. Are human hands and feet affected by climate? A test of Allen’s rule. *Am J Phys Anthropol* 158:132–140.
- Betti L, von Cramon-Taubadel N, Lycett SJ. 2012. Human pelvis and long bones reveal differential preservation of ancient population history and migration out of Africa. *Hum Biol* 84:139–152.
- Betti L, von Cramon-Taubadel N, Manica A, Lycett SJ. 2013. Global geometric morphometric analyses of the human pelvis reveal substantial neutral population history effects, even across sexes. *PLOS One* 8:e55909.
- Betti L, von Cramon-Taubadel N, Manica A, Lycett SJ. 2014. The interaction of neutral evolutionary processes with climatically driven adaptive changes in the 3D shape of the human os coxae. *J Hum Evol* 73:64–74.
- Bogin B, Rios L. 2003. Rapid morphological change in living humans: implications for modern human origins. *Comp Biochem Phys A* 136:71–84.
- Bogin B, Smith P, Orden AB, Varela Silva MI, Loucky J. 2002. Rapid change in height and body proportions of Maya American children. *Am J Hum Biol* 14:753–761.
- Bramble DM, Lieberman DE. 2004. Endurance running and the evolution of *Homo*. *Nature* 432:345–352.
- Brothwell DR. 1981. Digging up bones. Oxford: Oxford University Press.
- Caldwell W, Moloy H. 1933. Anatomical variations in the female pelvis and their effect in labor with a suggested classification. *Am J Obstet Gynecol* 26:479–505.
- Caldwell W, Moloy H. 1938. Anatomical variations in the female pelvis: their classification and obstetrical significance. *Proc R Soc Med* 32:1–30.
- Cavalli-Sforza LL, Menozzi P, Piazza A. 1994. The history and geography of human genes. Princeton: Princeton University Press.
- Cnattingius R, Cnattingius S, Notzon FC. 1998. Obstacles to reducing cesarean rates in a low-cesarean setting: the effect of maternal age, height, and weight. *Obstet Gynecol* 92:501–506.
- Coon CS. 1962. The origin of races. New York: Alfred A. Knopf.
- Cottingham J, Royston E. 1991. Obstetric fistulae: a review of available information. Geneva: World Health Organization.
- Crognier E. 1981. Climate and anthropometric variations in Europe and the Mediterranean area. *Ann Hum Biol* 8:99–107.
- Derry DE. 1923. On the sexual and racial characters of the human ilium. *J Anat* 58:71–83.
- DeSilva JM. 2011. A shift toward birthing relatively large infants early in human evolution. *Proc Natl Acad Sci USA* 108:1022–1027.
- Dolea C, AbouZhar C. 2003. Global burden of obstructed labour in the year 2000. Geneva, Switzerland: World Health Organisation.
- Dunsworth HM, Warrener AG, Deacon T, Ellison PT, Pontzer H. 2012. Metabolic hypothesis for human altriciality. *Proc Natl Acad Sci USA* 109:15212–15216.
- Eller E. 1999. Population substructure and isolation by distance in three continental regions. *Am J Phys Anthropol* 108:147–159.
- Eriksson A, Betti L, Friend AD, Lycett SJ, Singarayer JS, von Cramon-Taubadel N, Valdes PJ, Balloux F, Manica A. 2012. Late Pleistocene climate change and the global expansion of anatomically modern humans. *Proc Natl Acad Sci USA* 109:16089–16094.
- Fischer B, Mitteroecker P. 2015. Covariation between human pelvis shape, stature, and head size alleviates the obstetric dilemma. *Proc Natl Acad Sci USA* 112:5655–5660.
- Fu Q, Mittnik A, Johnson PL, Bos K, Lai M, Bollongino R, Sun C, Giemsch L, Schmitz R, Burger J, Ronchitelli AM, Martini F, Cremonesi RG, Svoboda J, Bauer P, Caramelli D, Castellano S, Reich D, Pääbo S, Krause J. 2013. A revised timescale for human evolution based on ancient mitochondrial genomes. *Curr Biol* 23:553–559.



- Grabowski MW. 2013. Hominin obstetrics and the evolution of constraints. *Evol Biol* 40:57–75.
- Grabowski MW. 2016. Bigger brains led to bigger bodies? The correlated evolution of human brain and body size. *Curr Anthropol* 57:174–196.
- Grabowski MW, Polk JD, Roseman CC. 2011. Divergent patterns of integration and reduced constraint in the human hip and the origins of bipedalism. *Evolution* 65:1336–1356.
- Grabowski M, Roseman CC. 2015. Complex and changing patterns of natural selection explain the evolution of the human hip. *J Hum Evol* 85:94–110.
- Greulich WW, Thoms H. 1938. The dimensions of the pelvic inlet of 789 white females. *Anat Rec* 72:45–51.
- Greulich WW, Thoms H, Twaddle R. 1939. A study of pelvic type and its relationship to body build in white women. *J Am Med Assoc* 112:485–493.
- Hanihara T. 2008. Morphological variation of major human populations based on nonmetric dental traits. *Am J Phys Anthropol* 136:169–182.
- Harvati K, Weaver TD. 2006. Human cranial anatomy and the differential preservation of population history and climate signatures. *Anat Rec* 288A:1225–1233.
- Hogan MC, Foreman KJ, Naghavi M, Ahn SY, Wang M, Makela SM, Lopez AD, Lozano R, Murray CJ. 2010. Maternal mortality for 181 countries, 1980–2008: a systematic analysis of progress towards Millennium Development Goal 5. *Lancet* 375:1609–1623.
- Holland EL, Cran GW, Elwood JH, Pinkerton JHM, Thompson W. 1982. Associations between pelvic anatomy, height and year of birth of men and women in Belfast. *Annals of Hum Biol* 9:113–120.
- Holliday TW. 1997a. Body proportions in Late Pleistocene Europe and modern human origins. *J Hum Evol* 32:423–447.
- Holliday TW. 1997b. Postcranial evidence of cold adaptation in European Neandertals. *Am J Phys Anthropol* 104:245–258.
- Holliday TW. 1999. Brachial and crural indices of European late upper paleolithic and mesolithic humans. *J Hum Evol* 36:549–566.
- Holliday TW. 2002. Body size and postcranial robusticity of European upper paleolithic hominins. *J Hum Evol* 43:513–528.
- Holliday TW, Falsetti AB. 1995. Lower limb length of European early modern humans in relation to mobility and climate. *J Hum Evol* 29:141–153.
- Holliday TW, Hilton CE. 2010. Body proportions of circumpolar peoples as evidenced from skeletal data: Ipiutak and Tigara (Point Hope) versus Kodiak Island Inuit. *Am J Phys Anthropol* 142:287–302.
- Hubbe M, Hanihara T, Harvati K. 2009. Climate signatures in the morphological differentiation of worldwide modern human populations. *Anat Rec* 292:1720–1733.
- Huseynov A, Zollikofer CPE, Coudyzer W, Gascho D, Kellenberger C, Hinzpeter R, Ponce de León MS. 2016. Developmental evidence for obstetric adaptation of the human female pelvis. *Proc Natl Acad Sci USA* 113:5227–5232.
- Imaizumi Y, Morton NE, Lalouel JM. 1973. Kinship and race. In: Morton NE, editor. *Genetic structure of populations*. Honolulu: University Press of Hawaii. p 228–233.
- Işcan MY. 1983. Assessment of race from the pelvis. *Am J Phys Anthropol* 62:205–208.
- Katzmarzyk PT, Leonard WR. 1998. Climatic influences on human body size and proportions: ecological adaptations and secular trends. *Am J Phys Anthropol* 106:483–503.
- Kelley MA. 1978. Phenice's visual sexing technique for the os pubis: a critique. *Am J Phys Anthropol* 48:121–122.
- Kelley JO, Angel JL. 1987. Life stresses of slavery. *Am J Phys Anthropol* 74:199–211.
- Kolawole TM, Adamu SP, Evans KT. 1978. Comparative pelvimetric measurements in Nigerian and Welsh women. *Clin Radiol* 29:85–90.
- Kriewall TJ, McPherson GK. 1981. Effects of uterine contractility on the fetal cranium: perspectives from the past, present and future. In: Milunsky A, Friedman EA, Gluck L, editors. *Advances in perinatal medicine*, vol. 1. New York: Plenum Medical Book Company. p 295–356.
- Krukier S. 1951. The sexual differences of the human pelvis. *Gynaecologia* 132:92–110.
- Krukier S. 1954. The problem of the normal pelvis. *Gynaecologia* 137:1–10.
- Kurki HK. 2007. Protection of obstetric dimensions in a small-bodied human sample. *Am J Phys Anthropol* 133:1152–1165.
- Kurki HK. 2013a. Bony pelvic canal size and shape in relation to body proportionality in humans. *Am J Phys Anthropol* 151:88–101.
- Kurki HK. 2013b. Skeletal variability in the pelvis and limb skeleton of humans: does stabilizing selection limit female pelvic variation? *Am J Hum Biol* 25:796–802.
- Kwawukume EY, Ghosh TS, Wilson JB. 1993. Maternal height as a predictor of vaginal delivery. *Int J Gynaecol Obstet* 41:27–30.
- LaVelle M. 1995. Natural selection and developmental sexual variation in the human pelvis. *Am J Phys Anthropol* 98:59–72.
- Leary S, Fall C, Osmond C, Lovel H, Campbell D, Eriksson J, Forrester T, Godfrey K, Hill J, Jie M, Law C, Newby R, Robinson S, Yajnik CS. 2006. Geographical variation in relationships between parental body size and offspring phenotype at birth. *Acta Obstet Gynecol Scand* 85:1066–1079.
- Leutenegger W. 1974. Functional aspects of pelvic morphology in simian Primates. *J Hum Evol* 3:207–222.
- Lovejoy CO. 1988. Evolution of human walking. *Sci Am* 259:118–125.
- Lovejoy CO. 2005. The natural history of human gait and posture. Part 1. Spine and pelvis. *Gait Posture* 21:95–112.
- Lovejoy CO, Cohn MJ, White TD. 1999. Morphological analysis of the mammalian postcranium: a developmental perspective. *Proc Natl Acad Sci USA* 96:13247–13252.
- Lovejoy CO, Heiple KG, Burstein AH. 1973. The gait of Australopithecus. *Am J Phys Anthropol* 38:757–779.
- Manica A, Amos W, Balloux F, Hanihara T. 2007. The effect of ancient population bottlenecks on human phenotypic variation. *Nature* 448:346–U346.
- McDougall I, Brown FH, Fleagle JG. 2005. Stratigraphic placement and age of modern humans from Kibish, Ethiopia. *Nature* 433:733–736.
- Mellars P. 2006. Why did modern human populations disperse from Africa ca. 60,000 years ago? A new model. *Proc Natl Acad Sci USA* 103:9381–9386.
- Micozzi MS. 1982. Skeletal tuberculosis, pelvic contraction, and parturition. *Am J Phys Anthropol* 58:441–445.
- Newman MT. 1953. The application of ecological rules to the racial anthropology of the Aboriginal New World. *Am Anthropol* 55:311–327.
- Newman MT. 1961. Biological adaptation of man to his environment: heat, cold, altitude, and nutrition. *Ann N Y Acad Sci* 91:617–633.
- Nicholson C. 1946. Variation in the female pelvis. *Lancet* 248:192–195.
- Oxorn H. 1986. *Oxorn-Foote human labor and birth*, fifth edition. Norwalk, Connecticut: Appleton-Century-Croft.
- Patriquin ML, Steyn M, Loth SR. 2002. Metric assessment of race from the pelvis in South Africans. *Forensic Sci Int* 127:104–113.
- Pearson OM. 2000. Activity, climate, and postcranial robusticity: implications for modern human origins and scenarios of adaptive change. *Curr Anthropol* 41:569–607.
- Ponce de León MS, Golovanova L, Doronichev V, Romanova G, Akazawa T, Kondo O, Ishida H, Zollikofer CPE. 2008. Neanderthal brain size at birth provides insights into the evolution of human life history. *Proc Natl Acad Sci USA* 105:13764–13768.
- Prasad M, Al-Tajer H. 2002. Maternal height and labour outcome. *J Obstet Gynaecol* 22:513–515.
- Prugnolle F, Manica A, Balloux F. 2005. Geography predicts neutral genetic diversity of human populations. *Curr Biol* 15:R159–R160.
- Ramachandran S, Deshpande O, Roseman CC, Rosenberg NA, Feldman MW, Cavalli-Sforza LL. 2005. Support from the relationship of genetic and geographic distance in human populations for

- a serial founder effect originating in Africa. *Proc Natl Acad Sci USA* 102:15942–15947.
- Relethford JH. 1994. Craniometric variation among modern human populations. *Am J Phys Anthropol* 95:53–62.
- Relethford JH. 2002. Apportionment of global human genetic diversity based on craniometrics and skin color. *Am J Phys Anthropol* 118:393–398.
- Relethford JH. 2004a. Boas and beyond: migration and craniometric variation. *Am J Hum Biol* 16:379–386.
- Relethford JH. 2004b. Global patterns of isolation by distance based on genetic and morphological data. *Hum Biol* 76:499–513.
- Rightmire GP. 2004. Brain size and encephalization in early to Mid-Pleistocene Homo. *Am J Phys Anthropol* 124:109–123.
- Roberts DF. 1953. Body weight, race and climate. *Am J Phys Anthropol* 11:533–558.
- Roberts DF. 1978. *Climate and human variability*. Menlo Park: Cummings.
- Rolian C. 2014. Genes, development, and evolvability in primate evolution. *Evol Anthropol* 23:93–104.
- Roseman CC. 2004. Detecting interregionally diversifying natural selection on modern human cranial form by using matched molecular and morphometric data. *Proc Natl Acad Sci USA* 101:12824–12829.
- Roseman CC, Auerbach BM. 2015. Ecogeography, genetics, and the evolution of human body form. *J Hum Evol* 78:80–90.
- Roseman CC, Weaver TD. 2004. Multivariate apportionment of global human craniometric diversity. *Am J Phys Anthropol* 125:257–263.
- Rosenberg K, Trevathan W. 1995. Bipedalism and human birth: the obstetrical dilemma revisited. *Evol Anthropol* 4:161–168.
- Rosenberg K, Trevathan W. 2002. Birth, obstetrics and human evolution. *Brit J Obstet Gynaecol* 109:1199–1206.
- Rosenberg KR. 1992. The evolution of modern human childbirth. *Am J Phys Anthropol* 35:89–124.
- Rosenberg KR, Brace CL, Frayer DW, Geise MC, Greene DL, Tague RG, Tappen NC, Trevathan WR, Trinkaus E. 1988. The functional significance of Neandertal pubic length. *Curr Anthropol* 29:595–617.
- Ruff C. 1991. Climate and body shape in hominid evolution. *J Hum Evol* 21:81–105.
- Ruff C. 1993. Climatic adaptation and hominid evolution: the thermoregulatory imperative. *Evol Anthropol* 2:53–60.
- Ruff C. 1994. Morphological adaptation to climate in modern and fossil hominids. *Am J Phys Anthropol* 37:65–107.
- Ruff C. 2002. Variation in human body size and shape. *Annu Rev Anthropol* 31:211–232.
- Ruff CB, Trinkaus E, Holliday TW. 1997. Body mass and encephalization in Pleistocene Homo. *Nature* 387:173–176.
- Schreider E. 1950. Geographical distribution of the body-weight/body-surface ratio. *Nature* 165:286–286.
- Schreider E. 1964. Ecological rules, body-heat regulation, and human evolution. *Evolution* 18:1–9.
- Schreider E. 1975. Morphological variations and climatic differences. *J Hum Evol* 4:529–539.
- Schultz AH. 1949. Sex differences in the pelvis of primates. *Am J Phys Anthropol* 7:401–424.
- Schutz H, Donovan ER, Hayes JP. 2009. Effects of parity on pelvic size and shape dimorphism in *Mus*. *J Morphol* 270:834–842.
- Sheiner E, Levy A, Katz M, Mazar M. 2005. Short-stature—an independent risk factor for Cesarean delivery. *Eur J Obstet Gynecol Reprod Biol* 120:175–178.
- Sibley LM, Armelagos GJ, Van Gerven DP. 1992. Obstetric dimensions of the true pelvis in a Medieval population from Sudanese Nubia. *Am J Phys Anthropol* 89:421–430.
- Smith HF. 2009. Which cranial regions reflect molecular distances reliably in humans? Evidence from three-dimensional morphology. *Am J Hum Biol* 21:36–47.
- Smith HF. 2011. The role of genetic drift in shaping modern human cranial evolution: a test using microevolutionary modeling. *Int J Evol Biol* 2011:145262.
- Specker B, Binkley T. 2005. High parity is associated with increased bone size and strength. *Osteoporosis Int* 16:1969–1974.
- Stewart KS, Philpott RH. 1980. Fetal response to cephalopelvic disproportion. *Br J Obstet Gynaecol* 87:641–649.
- Stock JT. 2006. Hunter-gatherer postcranial robusticity relative to patterns of mobility, climatic adaptation, and selection for tissue economy. *Am J Phys Anthropol* 131:194–204.
- Straus WL Jr. 1927. The human ilium: sex and stock. *Am J Phys Anthropol* 11:1–28.
- Strauss A, Hubbe M. 2010. Craniometric similarities within and between human populations in comparison with neutral genetic data. *Hum Biol* 82:315–330.
- Stuart-Macadam PL. 1989. Nutritional deficiency diseases: a survey of scurvy, rickets, and iron-deficiency anemia. In: Işcan MY, Kennedy KAR, editors. *Reconstruction of life from the skeleton*. New York: Alan R. Liss. p 201–222.
- Tague RG. 1989. Variation in pelvic size between males and females. *Am J Phys Anthropol* 80:59–71.
- Tague RG. 1995. Variation in pelvic size between males and females in nonhuman anthropoids. *Am J Phys Anthropol* 97:213–233.
- Tague RG, Lovejoy CO. 1986. The obstetric pelvis of A.L. 288-1 (Lucy). *J Hum Evol* 15:237–255.
- Thoms H. 1946. A discussion of pelvic variation and a report on the findings in 100 Negro women. *Am J Obstet Gynecol* 52:248–254.
- Thoms H. 1947. The role of nutrition in pelvic variation. *Am J Obstet Gynecol* 54:62–67.
- Tilkens MJ, Wall-Scheffler C, Weaver TD, Steudel-Numbers K. 2007. The effects of body proportions on thermoregulation: an experimental assessment of Allen's rule. *J Hum Evol* 53:286–291.
- Trevathan W, Rosenberg K. 2000. The shoulders follow the head: postcranial constraints on human childbirth. *J Hum Evol* 39:583–586.
- Trinkaus E. 1981. Neanderthal limb proportions and cold adaptation. In: Stringer CB, editor. *Aspects of human evolution*. London: Taylor and Francis. p 187–224.
- Tsu VD. 1992. Maternal height and age: risk factors for cephalopelvic disproportion in Zimbabwe. *Int J Epidemiol* 21:941–947.
- Turner WM. 1885. The index of the pelvic brim as a basis of classification. *J Anat Physiol* 20:125–143.
- von Cramon-Taubadel N. 2009a. Congruence of individual cranial bone morphology and neutral molecular affinity patterns in modern humans. *Am J Phys Anthropol* 140:205–215.
- von Cramon-Taubadel N. 2009b. Revisiting the homology hypothesis: the impact of phenotypic plasticity on the reconstruction of human population history from craniometric data. *J Hum Evol* 57:179–190.
- von Cramon-Taubadel N. 2011. The relative efficacy of functional and developmental cranial modules for reconstructing global human population history. *Am J Phys Anthropol* 146:83–93.
- von Cramon-Taubadel N, Lycett SJ. 2008. Brief communication: human cranial variation fits iterative founder effect model with African origin. *Am J Phys Anthropol* 136:108–113.
- von Cramon-Taubadel N, Weaver TD. 2009. Insights from a quantitative genetic approach to human morphological evolution. *Evol Anthropol* 18:237–240.
- Wall-Scheffler CM. 2012. Energetics, locomotion, and female reproduction: implications for human evolution. *Annu Rev Anthropol* 41:71–85.
- Warrener AG, Lewton KL, Pontzer H, Lieberman DE. 2015. A wider pelvis does not increase locomotor cost in humans, with implications for the evolution of childbirth. *PLOS One* 10:e0118903.
- Washburn SN. 1960. Tools and human evolution. *SciAm* 203:3–15.
- Weaver TD. 2002. A multi-causal functional analysis of hominid hip morphology. Palo Alto: Stanford University.
- Weaver TD, Hublin JJ. 2009. Neandertal birth canal shape and the evolution of human childbirth. *Proc Natl Acad Sci USA* 106:8151–8156.
- Weaver TD, Roseman CC, Stringer CB. 2007. Were Neandertal and modern human cranial differences produced by natural selection or genetic drift? *J Hum Evol* 53:135–145.
- Weaver TD, Roseman CC, Stringer CB. 2008. Close correspondence between quantitative- and molecular-genetic divergence times for Neandertals and modern humans. *Proc Natl Acad Sci USA* 105:4645–4649.

- Weaver TD, Steudel-Numbers K. 2005. Does climate or mobility explain the differences in body proportions between Neandertals and their Upper Paleolithic successors?. *Evol Anthropol* 14:218–223.
- Weinstein KJ. 2005. Body proportions in ancient Andeans from high and low altitudes. *Am J Phys Anthropol* 128:569–585.
- Wells JCK, DeSilva JM, Stock JT. 2012. The obstetric dilemma: an ancient game of Russian roulette, or a variable dilemma sensitive to ecology? *Yearb Phys Anthropol* 55:40–71.
- White TD, Black MT, Folkens PA. 2012. *Human osteology*. San Diego: Academic Press.
- Wilson LAB, Ives R, Cardoso HFV, Humphrey LT. 2015. Shape, size, and maturity trajectories of the human ilium. *Am J Phys Anthropol* 156:19–34.
- Witter FR, Caulfield LE, Stoltzfus RJ. 1995. Influence of maternal anthropometric status and birth weight on the risk of caesarean delivery. *Obstet Gynecol* 85:947–951.
- Wittman AB, Wall LL. 2007. The evolutionary origins of obstructed labor: bipedalism, encephalization, and the human obstetric dilemma. *Obstet Gynecol Surv* 62:739–748.