

A novel reduction of alcohols and ethers with a HSiEt₃/catalytic B(C₆F₅)₃ system

Vladimir Gevorgyan,^{a,*} Jian-Xiu Liu,^b Michael Rubin,^a Sharonda Benson^a and
Yoshinori Yamamoto^{b,*}

^aDepartment of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607-7061, USA


^bDepartment of Chemistry, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan

Received 29 July 1999; accepted 14 September 1999

Abstract













The primary alcohols **1a–d** and ethers **4a–b** were effectively reduced into the corresponding hydrocarbons **2** by HSiEt₃ in the presence of catalytic amounts of B(C₆F₅)₃. The secondary alkyl ethers **4g,h** underwent cleavage and/or reduction under similar reaction conditions to produce either the silyl ether **3k** or the corresponding alcohol **5b** upon subsequent deprotection with TBAF. The secondary alcohols (**1g,h**) and tertiary alcohol **1i**, as well as tertiary alkyl ether **4i**, did not react with the HSiEt₃/B(C₆F₅)₃ reducing reagent at all. The following relative reactivity order of substrates was found: primary>>secondary>tertiary. The methyl aryl ethers **4c–e** and alkyl aryl ether **4f** were smoothly deprotected to give the corresponding silyl ethers **3b,h–j** in nearly quantitative isolated yields. © 1999 Elsevier Science Ltd. All rights reserved.

It is known that hydrosilanes are able to reduce alcohols and alkyl ethers in the presence of Lewis acids (Eq. 1).¹ These methods require at least stoichiometric amounts of Lewis acid. Furthermore, the methodologies described are most effective for the reduction of a C–O bond at tertiary carbon,¹ are much less effective for the reduction of secondary substrates,² and absolutely ineffective for the reduction of primary alcohols and ethers (Eq. 1).¹



* Corresponding authors. Tel: +81 22 217 6581; fax: +81 22 217 6784; e-mail: yoshi@yamamoto1.chem.tohoku.ac.jp
Tel: +1 312 355 3579; fax: +1 312 355 0836; e-mail: vald@tiger.cc.uic.edu

Table 1
Reduction of alcohols **1** with HSiEt₃/cat.-B(C₆H₅)₃ system^a

















Entry	Alcohol 1	HSiEt ₃ (eq)	Products (Yield, %)
1	<i>n</i> -C ₉ H ₁₉ OH (1a)	1.1	<i>n</i> -C ₉ H ₁₉ OSiEt ₃ (3a) (>95) ^b
2	(1a)	3.0	<i>n</i> -C ₉ H ₂₀ (2a) (>99) ^c
3	 (1b)	6.0	 (2b) (91) ^c
4	Ph(CH ₂) ₃ OH (1c)	3.0	Ph(CH ₂) ₃ H (2c) (>95) ^b
5	Ph(CH ₂) ₂ OH (1d)	3.0	Ph(CH ₂) ₂ H (2d) (>95) ^b
6	 (1e)	1.1 (or 6.0)	 (3b) (>99) ^c
7	 (1f)	3.0	 (3c) (>95) ^b
8	 (1g)	6.0	 (3d) (>95) ^b
9	 (1h)	6.0	 (3e) (>95) ^b
	 (1i)		 (3f) (86) ^b
10		3.0	

^aHSiEt₃ was added under an argon atmosphere to a mixture of B(C₆F₅)₃ (10 mol %) and alcohol **1** (1 mmol) in hexane or CH₂Cl₂ (0.5 mL). After being stirred for 20 hours at room temperature, the appropriate internal standard was added and the mixture was analyzed by capillary GC. For NMR analysis, after completion of the reaction the mixture was quenched (Et₃N, 0.5 mL), filtered through celite and concentrated, followed by addition of internal standard (CH₃CCl₃). In the case of low b.p. products the reaction mixture was analyzed by ¹H NMR without concentration. ^bNMR yield. ^cGC yield.

We found that HSiEt₃ (3.0 equiv.) in the presence of catalytic amounts of B(C₆F₅)₃ (10 mol%) enabled us to reduce primary alcohols **1a–d** into the corresponding hydrocarbons **2a–d** in very high to quantitative yields (Eq. 2, Table 1, entries 2–5). It should be noted that employment of an equimolar amount of HSiEt₃ was not sufficient for the reduction of **1a**, instead alkyl silyl ether **3a** was produced quantitatively (entry 1).³ In contrast to the primary alcohols, the more bulky secondary alcohols **1f–h**, as well as the tertiary alcohol **1i**, were stable towards reduction, and the corresponding silyl ethers **3c–f** were obtained in very high yields (entries 7–10). Phenol, as expected, did not undergo reduction even under treatment with a large excess of HSiEt₃ (entry 6).



Table 2
Cleavage and reduction of ethers **4** with HSiEt₃/cat.-B(C₆H₅)₃ system^a


Entry	Ether 4	HSiEt ₃ (eq)	Products (Yield, %)
1	(<i>n</i> -C ₁₀ H ₂₁) ₂ O (4a)	1.1	<i>n</i> -C ₁₀ H ₂₂ (2e) (93) ^b <i>n</i> -C ₁₀ H ₂₁ OSiEt ₃ (3g) (95) ^c
2	(4a)	3.0	2 × (2e) (>99) ^b
3	 (4b)	1.1	 (5a) (88) ^{c,d}
4	(4b)	3.0	 (2f) (87) ^c
5	 (4c)	1.1	 (3b) (>99) ^{e,f}
6	 (4d)	1.1	 (3h) (96) ^e
7	 (4e)	1.1	 (3i) (98) ^e
8	 (4f)	1.1	 (3j) (>99) ^{e,f}
9	 (4g)	3.0	 (3k) (>95) ^c
10	 (4h)	3.0	 (5b) (91) ^{c,d}
11	 (4i)		no reaction ^g

^a All reactions were carried in CH₂Cl₂. For more detailed reaction conditions, see Table 1, note a. ^bGC yield. ^cNMR yield. ^dYields of alcohols **5** after TBAF deprotection of corresponding alkoxysilanes **3**. ^eIsolated yield. ^fReaction was performed in 5 mmol scale.



^g97% of **4a** was recovered.

Next, we tested the HSiEt₃/cat.-B(C₆F₅)₃ reducing system for cleavage of alkyl ethers. It was found that linear primary alkyl ether **4a** could be easily cleaved in the presence of a stoichiometric amount of HSiEt₃/10 mol% B(C₆F₅)₃ to give decane (**2e**) and silylether **3g** (Eq. 3, Table 2, entry 1). Similarly, reduction of cyclic primary ether **4b** gave the corresponding ring cleaved silyl ether, which was converted

into alcohol **5a** in 88% yield upon treatment with TBAF (entry 3). However, in the presence of excess amounts of HSiEt_3 both linear and cyclic ethers **4a,b** underwent smooth exhaustive reduction into the hydrocarbons **2e,f** in quantitative to high yields, respectively (entries 2 and 4). It was found that methyl aryl ethers **4c–e** readily underwent deprotection under the treatment of 1.1 equiv. of HSiEt_3 to give the corresponding silyl ethers **3b,h,i** in virtually quantitative yields (Table 2, entries 5–7). As expected, the aryl C–O bond in dehydrobenzofuran (**4e**) was tolerant towards the reduction, and consequently the cleavage product **3j** was produced quantitatively (entry 8). The secondary alkyl ether **4g** was quantitatively cleaved in the presence of 3 equivalents of HSiEt_3 to give the silyl ether **3k** (entry 9), thus exhibiting a striking resistance of linear secondary alkyl ethers (as well as corresponding alcohols; see also Table 1, entries 7–9) towards reduction. Cyclic secondary ether **4h** behaved similarly, producing the cleavage product **5b** in very high yield (entry 10). Sterically hindered ether **4i**, possessing both tertiary and primary alkyl units, did not undergo the reduction at all (entry 11).



In conclusion, we developed a novel effective method for reduction of primary alcohols and ethers with HSiEt_3 in the presence of catalytic amounts of $\text{B(C}_6\text{F}_5)_3$ (Eq. 4). In addition, this method could serve as a powerful tool for deprotection of methyl aryl and alkyl aryl ethers. Further development of this synthetic methodology, as well as mechanistic studies, are now underway in our laboratories.



References

1. (a) Adlington, M. G.; Orfanopoulos, M.; Fry, J. L. *Tetrahedron Lett.* **1976**, 2955. (b) Fry, J. L.; Orfanopolous, M.; Adlington, M. G.; Dittman, W. R.; Silverman, S. B. *J. Org. Chem.* **1978**, *43*, 374. (c) Orfanopolous, M.; Smonou, I. *Synthetic Commun.* **1988**, *18*, 833. (d) Larsen, J. W.; Chang, L. W. *J. Org. Chem.* **1979**, *44*, 1168. (e) Yato, M.; Ishida, A. *Heterocycles* **1995**, *41*, 17. (f) Smonou, I. *Synthetic Commun.* **1994**, *24*, 1999.
2. For a report on the reduction of secondary benzyl alcohols in the presence of primary alkyl alcohols with $\text{HSiEt}_3/\text{BF}_3$ system, see Ref. 1c.
3. During the preparation of this manuscript, a method for silyl protection of alcohols with hydrosilanes in the presence of $\text{B(C}_6\text{F}_5)_3$ was reported, see: Blackwell, J. M.; Foster, K. L.; Beck, V. H.; Piers, W. E. *J. Org. Chem.* **1999**, *64*, 4887.